

## PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>7</sup> : <b>A61K 31/00</b></p>	<p><b>A2</b></p>	<p>(11) International Publication Number: <b>WO 00/50020</b></p> <p>(43) International Publication Date: <b>31 August 2000 (31.08.00)</b></p>								
<table border="0" style="width: 100%;"> <tr> <td style="width: 50%; vertical-align: top;"> <p>(21) International Application Number: <b>PCT/US00/04187</b></p> <p>(22) International Filing Date: <b>18 February 2000 (18.02.00)</b></p> <p>(30) Priority Data:</p> <table border="0" style="width: 100%;"> <tr> <td style="width: 33%;">60/121,339</td> <td style="width: 33%;">24 February 1999 (24.02.99)</td> <td style="width: 33%;">US</td> </tr> <tr> <td>Not furnished</td> <td>18 February 2000 (18.02.00)</td> <td>US</td> </tr> </table> <p>(71) Applicant (for all designated States except US): <b>UNIVERSITY OF CINCINNATI [US/US];</b> Wherry Hall G-07, Box 670829, 3223 Eden Avenue, Cincinnati, OH 45267-0829 (US).</p> <p>(72) Inventor; and</p> <p>(75) Inventor/Applicant (for US only): <b>MCELROY, Susan, L. [US/US];</b> 8395 Old Stable Road, Cincinnati, OH 45243-1441 (US).</p> <p>(74) Agents: <b>GOLDSSTEIN, Steven, J. et al.; Frost &amp; Jacobs LLP,</b> 2500 PNC Center, 201 E. Fifth Street, Cincinnati, OH 45202 (US).</p> </td> <td style="width: 50%; vertical-align: top;"> <p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> Without international search report and to be republished upon receipt of that report.</p> </td> </tr> </table>			<p>(21) International Application Number: <b>PCT/US00/04187</b></p> <p>(22) International Filing Date: <b>18 February 2000 (18.02.00)</b></p> <p>(30) Priority Data:</p> <table border="0" style="width: 100%;"> <tr> <td style="width: 33%;">60/121,339</td> <td style="width: 33%;">24 February 1999 (24.02.99)</td> <td style="width: 33%;">US</td> </tr> <tr> <td>Not furnished</td> <td>18 February 2000 (18.02.00)</td> <td>US</td> </tr> </table> <p>(71) Applicant (for all designated States except US): <b>UNIVERSITY OF CINCINNATI [US/US];</b> Wherry Hall G-07, Box 670829, 3223 Eden Avenue, Cincinnati, OH 45267-0829 (US).</p> <p>(72) Inventor; and</p> <p>(75) Inventor/Applicant (for US only): <b>MCELROY, Susan, L. [US/US];</b> 8395 Old Stable Road, Cincinnati, OH 45243-1441 (US).</p> <p>(74) Agents: <b>GOLDSSTEIN, Steven, J. et al.; Frost &amp; Jacobs LLP,</b> 2500 PNC Center, 201 E. Fifth Street, Cincinnati, OH 45202 (US).</p>	60/121,339	24 February 1999 (24.02.99)	US	Not furnished	18 February 2000 (18.02.00)	US	<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> Without international search report and to be republished upon receipt of that report.</p>
<p>(21) International Application Number: <b>PCT/US00/04187</b></p> <p>(22) International Filing Date: <b>18 February 2000 (18.02.00)</b></p> <p>(30) Priority Data:</p> <table border="0" style="width: 100%;"> <tr> <td style="width: 33%;">60/121,339</td> <td style="width: 33%;">24 February 1999 (24.02.99)</td> <td style="width: 33%;">US</td> </tr> <tr> <td>Not furnished</td> <td>18 February 2000 (18.02.00)</td> <td>US</td> </tr> </table> <p>(71) Applicant (for all designated States except US): <b>UNIVERSITY OF CINCINNATI [US/US];</b> Wherry Hall G-07, Box 670829, 3223 Eden Avenue, Cincinnati, OH 45267-0829 (US).</p> <p>(72) Inventor; and</p> <p>(75) Inventor/Applicant (for US only): <b>MCELROY, Susan, L. [US/US];</b> 8395 Old Stable Road, Cincinnati, OH 45243-1441 (US).</p> <p>(74) Agents: <b>GOLDSSTEIN, Steven, J. et al.; Frost &amp; Jacobs LLP,</b> 2500 PNC Center, 201 E. Fifth Street, Cincinnati, OH 45202 (US).</p>	60/121,339	24 February 1999 (24.02.99)	US	Not furnished	18 February 2000 (18.02.00)	US	<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> Without international search report and to be republished upon receipt of that report.</p>			
60/121,339	24 February 1999 (24.02.99)	US								
Not furnished	18 February 2000 (18.02.00)	US								
<p>(54) Title: <b>USE OF SULFAMATE DERIVATIVES FOR TREATING IMPULSE CONTROL DISORDERS</b></p> <div style="text-align: center; margin: 20px 0;"> <p style="margin-left: 400px;">(I)</p> </div> <p>(57) Abstract</p> <p>Impulse Control Disorders (ICDs) are characterized by harmful behaviors performed in response to irresistible impulses. The essential feature of an ICD is the failure to resist an impulse, drive, or temptation and to perform an act that is harmful to the person or to others. The present invention comprises methods for the treatment or prevention of ICDs using a class of sulfamates of formula (I) wherein X is CH<sub>2</sub> or oxygen, and R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub> and R<sub>5</sub> are as herein defined. Further, pharmaceutical compositions containing a compound of formula (I) as well as methods for their use and intermediates form part of the present invention are also disclosed.</p>										

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## USE OF SULFAMATE DERIVATIVES FOR TREATING IMPULSE CONTROL DISORDERS

5

SUSAN L. McELROY

This application is based on U.S. Provisional Patent Application Serial No.  
60/121,339, McElroy, filed February 24, 1999.

10

### **Technical Field**

The present invention generally relates to the use of drugs for the treatment of mental disorders. More specifically, the invention describes methods for the treatment and prevention of Impulse Control Disorders (ICD's) by administering sulfamate derivatives.

15

### **Background of the Invention**

Sulfamate derivatives having useful pharmaceutical activity in the areas of epilepsy, glaucoma, peptic ulcers and male infertility are disclosed in U.S. Pat. Nos. 4,075,351, 4,513,006, 4,591,601, 4,792,569, and 5,760,007. One of these compounds  
20 2,3:4,5-bis-O-(1-methylethylidene)- beta -D-fructopyranose sulfamate known as topiramate has been demonstrated in clinical trials of human epilepsy to be effective as adjunctive therapy or as monotherapy in treating simple and complex partial seizures and secondarily generalized seizures and is currently marketed for the treatment of simple and complex partial seizure epilepsy with or without secondary generalized seizures.

Binge eating disorder (BED) is characterized by discrete periods of binge eating during which large amounts of food are consumed in a discrete period of time and a sense of control over eating is absent. Persons with bulimia nervosa have been reported to have electroencephalographic abnormalities and to display reduced binge eating in response to the anti-epileptic drug phenytoin. Also, in controlled trials in patients with epilepsy, topiramate was associated with suppression of appetite and weight loss unrelated to binge eating.

Binge eating disorder is a subset of a larger classification of mental disorders broadly defined as Impulse Control Disorders (ICDs) characterized by harmful behaviors performed in response to irresistible impulses. It has been suggested that ICDs may be related to obsessive-compulsive disorder or similarly, maybe forms of obsessive-compulsive disorders. It has also been hypothesized that ICDs may be related to mood disorder or may be forms of affective spectrum disorder, a hypothesized family of disorders sharing at least one common physiologic abnormality with major depression.

In the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV), the essential feature of an ICD is the failure to resist an impulse, drive, or temptation to perform an act that is harmful to the person or to others. For most ICDs, the individual feels an increasing sense of tension or arousal before committing the act, and then experiences pleasure, gratification, or release at the time of committing the act. After the act is performed, there may or may not be regret or guilt. ICDs are listed in a residual category, the ICDs Not Elsewhere Classified, which includes intermittent explosive disorder (IED), kleptomania, pathological gambling, pyromania, trichotillomania, and ICD not otherwise specified (NOS). Examples of ICDs NOS are compulsive buying or shopping, repetitive

self-mutilation, nonparaphilic sexual addictions, severe nail biting, compulsive skin picking, personality disorders with impulsive features, attention deficit/hyperactivity disorder, eating disorders characterized by binge eating, and substance use disorders.

5

### Summary of the Invention


It is an object of the present invention to describe the use of sulfamate derivatives for the treatment of Impulse Control Disorders.

The present invention comprises methods for the treatment or prevention of Impulse Control Disorders using the compounds of formula (I), pharmaceutical  
10 compositions containing one or more of the compounds of formula (I), or pharmaceutical compositions containing one or more of the compounds of formula (I) in addition to a safe and effective amount of one or more additional agents to treat related symptoms and conditions.

15

### Detailed Description of the Invention


The sulfamates of use in the present invention are of the following formula (I):



wherein

X is CH<sub>2</sub> or oxygen;

R<sub>1</sub> is hydrogen or alkyl; and R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub> and R<sub>5</sub> are independently hydrogen or lower alkyl and, when X is CH<sub>2</sub>, R<sub>4</sub> and R<sub>5</sub> may be alkene groups joined to form a benzene ring and,  
 5 when X is oxygen, R<sub>2</sub> and R<sub>3</sub> and/or R<sub>4</sub> and R<sub>5</sub> together may be a methylenedioxy group of the following formula (II):



wherein

R<sub>6</sub> and R<sub>7</sub> are the same or different and are hydrogen, lower alkyl or are alkyl and are  
 10 joined to form a cyclopentyl or cyclohexyl ring. R<sub>1</sub> in particular is hydrogen or alkyl of about 1 to 4 carbons, such as methyl, ethyl and iso-propyl. Alkyl throughout this specification includes straight and branched chain alkyl. Alkyl groups for R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub> and R<sub>7</sub> are of about 1 to 3 carbons and include methyl, ethyl, iso-propyl and n-propyl. When X is CH<sub>2</sub>, R<sub>4</sub> and R<sub>5</sub> may combine to form a benzene ring fused to the 6-membered  
 15 X-containing ring, *i.e.*, R<sub>4</sub> and R<sub>5</sub> are defined by the alkatrienyl group =C-CH=CH-CH=.

A particular group of compounds of formula (I) is that wherein X is oxygen and both R<sub>2</sub> and R<sub>3</sub> and R<sub>4</sub> and R<sub>5</sub> together are methylenedioxy groups of the formula (II), wherein R<sub>6</sub> and R<sub>7</sub> are both hydrogen, both alkyl or combine to form a spiro cyclopentyl or cyclohexyl ring, in particular where R<sub>6</sub> and R<sub>7</sub> are both alkyl such as methyl. A second  
 20 group of compounds is that wherein X is CH<sub>2</sub> and R<sub>4</sub> and R<sub>5</sub> are joined to form a benzene

ring. A third group of compounds of formula (I) is that wherein both  $R_2$  and  $R_3$  are hydrogen.

The compounds of formula I: may be made by the processes disclosed in U.S. Pat. Nos. 4,075,351, 4,513,006, 4,591,601, 4,792,569 , 5,242,942, 5,387,700, which are  
5 incorporated in their entirety herein by reference.

The compounds of formula I include the various individual isomers as well as the racemates thereof. For treating ICDs, a compound of formula (I) may be employed at a daily dosage in the range of about 15 to 1400 mg administered orally, for an average adult human.

10 To prepare the pharmaceutical compositions of this invention, one or more sulfamate compounds of formula (I) are intimately admixed with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques, which carrier may take a wide variety of forms depending on the form of preparation desired for administration, *e.g.*, oral, by suppository, or parenteral. In preparing the compositions in  
15 oral dosage form, any of the usual pharmaceutical media may be employed. Thus, for liquid oral preparations, such as for example, suspensions, elixirs and solutions, suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like; for solid oral preparations such as, for example, powders, capsules and tablets, suitable carriers and additives include starches,  
20 sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar coated or enteric coated by standard

techniques. Suppositories may be prepared, in which case cocoa butter could be used as the carrier. For parenterals, the carrier will usually comprise sterile water, though other ingredients, for example, for purposes such as aiding solubility or for preservation, may be included. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Topiramate is currently available for oral administration in round tablets containing 25 mg, 100 mg or 200 mg of active agent. The tablets contain the following inactive ingredients: lactose hydrous, pregelatinized starch, microcrystalline cellulose, sodium starch glycolate, magnesium stearate, purified water, carnauba wax, hydroxypropyl methylcellulose, titanium dioxide, polyethylene glycol, synthetic iron oxide, and polysorbate 80.

The pharmaceutical compositions herein will contain, per dosage unit, *e.g.*, tablet, capsule, powder injection, teaspoonful, suppository and the like from about 5 to about 1000 mg of the active ingredient.

The activity of the compounds of formula I in treating ICD's was first evidenced in clinical studies conducted to evaluate the efficacy of topiramate in treating mood disorders. Several patients who coincidentally had binge eating disorder reported that there was a marked reduction in their bingeing and a concurrent loss in weight.

Examples of specific compounds of formula (I) are:

- 2,3:4,5-bis-O-(1-methylethylidene)- beta -D-fructopyranose sulfamate
- 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose sulfamate;
- 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -L-fructopyranose sulfamate;
- 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose methylsulfamate;
- 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose butylsulfamate;



- 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose ethylsulfamate;  
2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose octylsulfamate;  
2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose 2-  
propenylsulfamate;
- 5 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose  
phenylmethylsulfamate;  
2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose  
cyclopropylsulfamate;  
2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose  
10 cyclobutylsulfamate;  
2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose (2,2,2-  
trifluoroethyl)sulfamate;  
2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose  
dimethylsulfamate;
- 15 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose diethylsulfamate;  
2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose azidosulfate;  
(S)-2,3-O-(1-methylethylidene)-4,5-O-sulfinyl- beta -D-fructopyranose sulfamate;  
(R)-2,3-O-(1-methylethylidene)-4,5-O-sulfinyl- beta -D-fructopyranose sulfamate;  
2,3-O-(1-ethylpropylidene)-4,5-O-sulfonyl- beta -D-fructopyranose sulfamate;
- 20 2,3-O-(1-methylethylidene)-4,5-O-[N-(4-methylbenzenesulfonyl)imidodisulfonyl]- beta  
-D-fructopyranose sulfamate;  
2,3-O-(1-methylethylidene)-4,5-O-[N-(4-methylbenzenesulfonyl)imidodisulfonyl]- beta  
-D-fructopyranose sulfamate;

2,3-O-(cyclohexylidene)-4,5-O-sulfonyl- beta -D-fructopyranose sulfamate;  
(S)-4,5-O-[N-(1,1-dimethylethoxycarbonyl)imidosulfinyl]-2,3-O-(1-methylethylidene)- beta -D-fructopyranose sulfamate;  
and the pharmaceutically acceptable salts thereof.

5

Included within the scope of this invention are the various individual anomers, diastereomers and enantiomers as well as mixtures thereof. Such compounds are included within the definition of formula (I). In addition, the compounds of this invention also include any pharmaceutically acceptable salts, for example: alkali metal salts, such as  
10 sodium and potassium; ammonium salts; monoalkylammonium salts; dialkylammonium salts; trialkylammonium salts; tetraalkylammonium salts; and tromethamine salts. Hydrates and other solvates of the compound of the formula (I) are included within the scope of this invention.

Pharmaceutically acceptable salts of the compounds of formula (I) can be  
15 prepared by reacting the sulfamate of formula (I) with the appropriate base and recovering the salt.

The sulfamate derivatives may be used in conjunction with one or more other drug compound and used according to the methods of the present invention so long as the pharmaceutical agent has a use that is also effective in treating ICD's and/or concurrent  
20 illnesses. Pharmaceutical agents include the following categories and specific examples. It is not intended that the category be limited by the specific examples. Those of ordinary skill in the art will be able to identify readily those pharmaceutical agents that have utility with the present invention. Those of ordinary skill in the art will recognize also numerous

other compounds that fall within the categories and that are useful according to the invention.

- Adrenergic: Adrenalone; Amidephrine Mesylate; Apraclonidine  
Hydrochloride; Brimonidine Tartrate; Dapiprazole Hydrochloride; Deterenol  
5 Hydrochloride; Dipivefrin; Dopamine Hydrochloride; Ephedrine Sulfate;  
Epinephrine; Epinephrine Bitartrate; Epinephryl Borate; Esproquin  
Hydrochloride; Etafedrine Hydrochloride; Hydroxyamphetamine Hydrobromide;  
Levonordefrin; Mephentermine Sulfate; Metaraminol Bitartrate; Metizoline  
Hydrochloride; Naphazoline Hydrochloride; Norepinephrine Bitartrate;  
10 Oxidopamine; Oxymetazoline Hydrochloride; Phenylephrine Hydrochloride;  
Phenylpropanolamine Hydrochloride; Phenylpropanolamine Polistirex;  
Prenalterol Hydrochloride; Propylhexedrine; Pseudoephedrine Hydrochloride;  
Tetrahydrozoline Hydrochloride; Tramazoline Hydrochloride; Xylometazoline  
Hydrochloride.
- 15 Adrenocortical steroid: Ciprocinonide; Desoxycorticosterone Acetate;  
Desoxycorticosterone Pivalate; Dexamethasone Acetate; Fludrocortisone Acetate;  
Flumoxonide; Hydrocortisone Hemisuccinate; Methylprednisolone  
Hemisuccinate; Naflocort; Procinonide; Timobesone Acetate; Tipredane.
- Adrenocortical suppressant: Aminoglutethimide; Trilostane.
- 20 Alcohol deterrent: Disulfiram.
- Aldosterone antagonist: Canrenoate Potassium; Canrenone; Dicirenone;  
Mexrenoate Potassium; Prorenoate Potassium; Spironolactone.

Amino acid: Alanine; Aspartic Acid; Cysteine Hydrochloride; Cystine; Histidine; Isoleucine; Leucine; Lysine; Lysine Acetate; Lysine Hydrochloride; Methionine; Phenylalanine; Proline; Serine; Threonine; Tryptophan; Tyrosine; Valine.

5                   Analeptic: Modafinil.

                  Analgesic: Acetaminophen; Alfentanil Hydrochloride; Aminobenzoate Potassium; Aminobenzoate Sodium; Anidoxime; Anileridine; Anileridine Hydrochloride; Anilopam Hydrochloride; Anirolac; Antipyrine; Aspirin; Benoxaprofen; Benzydamine Hydrochloride; Bicifadine Hydrochloride; 10           Brifentanil Hydrochloride; Bromadoline Maleate; Bromfenac Sodium; Buprenorphine Hydrochloride; Butacetin; Butixirate; Butorphanol; Butorphanol Tartrate; Carbamazepine; Carbaspirin Calcium; Carbiphen Hydrochloride; Carfentanil Citrate; Ciprofadol Succinate; Ciramadol; Ciramadol Hydrochloride; Clonixeril; Clonixin; Codeine; Codeine Phosphate; Codeine Sulfate; Conorphone 15           Hydrochloride; Cyclazocine; Dexoxadrol Hydrochloride; Dexpemedolac; Dezocine; Diflunisal; Dihydrocodeine Bitartrate; Dimefadane; Dipyrone; Doxpicomine Hydrochloride; Drinidene; Enadoline Hydrochloride; Epirizole; Ergotamine Tartrate; Ethoxazene Hydrochloride; Etofenamate; Eugenol; Fenoprofen; Fenoprofen Calcium; Fentanyl Citrate; Floctafenine; Flufenisal; 20           Flunixin; Flunixin Meglumine; Flupirtine Maleate; Fluproquazone; Fluradoline Hydrochloride; Flurbiprofen; Hydromorphone Hydrochloride; Ibufenac; Indoprofen; Ketazocine; Ketorfanol; Ketorolac Tromethamine; Letimide Hydrochloride; Levomethadyl Acetate; Levomethadyl Acetate Hydrochloride;

Levonantradol Hydrochloride; Levorphanol Tartrate; Lofemizole Hydrochloride;  
Lofentanil Oxalate; Lorcinadol; Lomoxicam; Magnesium Salicylate; Mefenamic  
Acid; Menabitan Hydrochloride; Meperidine Hydrochloride; Meptazinol  
Hydrochloride; Methadone Hydrochloride; Methadyl Acetate; Methopholine;  
5 Methotrimeprazine; Metkephamid Acetate; Mimbane Hydrochloride; Mirfentanil  
Hydrochloride; Molinazone; Morphine Sulfate; Moxazocine; Nabitan  
Hydrochloride; Nalbuphine Hydrochloride; Nalmexone Hydrochloride;  
Namoxyrate; Nantradol Hydrochloride; Naproxen; Naproxen Sodium; Naproxol;  
Nefopam Hydrochloride; Nexeridine Hydrochloride; Noracymethadol  
10 Hydrochloride; Ocfentanil Hydrochloride; Octazamide; Olvanil; Oxetorone  
Fumarate; Oxycodone; Oxycodone Hydrochloride; Oxycodone Terephthalate;  
Oxymorphone Hydrochloride; Pemedolac; Pentamorphone; Pentazocine;  
Pentazocine Hydrochloride; Pentazocine Lactate; Phenazopyridine  
Hydrochloride; Phenyramidol Hydrochloride; Picenadol Hydrochloride;  
15 Pinadoline; Pirfenidone; Piroxicam Olamine; Pravadoline Maleate; Prodilidine  
Hydrochloride; Profadol Hydrochloride; Propirarn Fumarate; Propoxyphene  
Hydrochloride; Propoxyphene Napsylate; Proxazole; Proxazole Citrate;  
Proxorphan Tartrate; Pyrroliphen Hydrochloride; Remifentanil Hydrochloride;  
Salcolex; Salethamide Maleate; Salicylamide; Salicylate Meglumine; Salsalate;  
20 Sodium Salicylate; Spiradoline Mesylate; Sufentanil; Sufentanil Citrate;  
Talmacetin; Talniflumate; Talosalate; Tazadolene Succinate; Tebufelone;  
Tetrydamine; Tifurac Sodium; Tilidine Hydrochloride; Tiopinac; Tonazocine  
Mesylate; Tramadol Hydrochloride; Trefentanil Hydrochloride; Trolamine;

Veradoline Hydrochloride; Verilopam Hydrochloride; Volazocine; Xorphanol Mesylate; Xylazine Hydrochloride; Zenazocine Mesylate; Zomepirac Sodium; Zucapsaicin.

Anorectic compounds including dexfenfluramine.

5           Anorexic: Aminorex; Amphetamine Hydrochloride; Clominorex; Clortermine Hydrochloride; Diethylpropion Hydrochloride; Fenfluramine Hydrochloride; Fenisorex; Fludorex; Fluminorex; Levamfetamine Succinate; Mazindol; Mefenorex Hydrochloride; Phenmetrazine Hydrochloride; Phentermine; Sibutramine Hydrochloride.

10           Anti-anxiety agent: Adatanserin Hydrochloride; Alpidem; Binospirone Mesylate; Bretazenil; Glemanserin; Ipsapirone Hydrochloride; Mirisetrone Maleate; Ocinaclone; Ondansetron Hydrochloride; Panadiplon; Pancopride; Pazinaclone; Serazapine Hydrochloride; Tandospirone Citrate; Zalospirone Hydrochloride.

15           Antidepressant: Adatanserin Hydrochloride; Adinazolam; Adinazolam Mesylate; Alaproclate; Aletamine Hydrochloride; Amedalin Hydrochloride; Amitriptyline Hydrochloride; Amoxapine; Aptazapine Maleate; Azaloxan Fumarate; Azepindole; Azipramine Hydrochloride; Bipenamol Hydrochloride; Bupropion Hydrochloride; Butacetin; Butriptyline Hydrochloride; Caroxazone;  
20           Cartazolate; Ciclazindol; Cidoxepin Hydrochloride; Cilobamine Mesylate; Clodazon Hydrochloride; Clomipramine Hydrochloride; Cotinine Fumarate; Cyclindole; Cypenamine Hydrochloride; Cyprolidol Hydrochloride; Cyproximide; Daledalin Tosylate; Dapoxetine Hydrochloride; Dazadrol Maleate;

Dazepinil Hydrochloride; Desipramine Hydrochloride; Dexamisole; Deximafen;  
Dibenzepin Hydrochloride; Dioxadrol Hydrochloride; Dothiepin Hydrochloride;  
Doxepin Hydrochloride; Duloxetine Hydrochloride; Eclanamine Maleate;  
Encyprate; Etoperidone Hydrochloride; Fantridone Hydrochloride; Fehmetozole  
5 Hydrochloride; Fenmetramide; Fezolamine Fumarate; Fluotracen Hydrochloride;  
Fluoxetine; Fluoxetine Hydrochloride; Fluparoxan Hydrochloride; Gamfexine;  
Guanoxyfen Sulfate; Imafen Hydrochloride; Imiloxan Hydrochloride; Imipramine  
Hydrochloride; Indeloxazine Hydrochloride; Intripityline Hydrochloride;  
Iprindole; Isocarboxazid; Ketipramine Fumarate; Lofepramine Hydrochloride;  
10 Lortalamine; Maprotiline; Maprotiline Hydrochloride; Melitracen Hydrochloride;  
Milacemide Hydrochloride; Minaprine Hydrochloride; Mirtazapine;  
Moclobemide; Modaline Sulfate; Napactadine Hydrochloride; Napamezole  
Hydrochloride; Nefazodone Hydrochloride; Nisoxetine; Nitrafudam  
Hydrochloride; Nomifensine Maleate; Nortriptyline Hydrochloride; Octriptyline  
15 Phosphate; Opipramol Hydrochloride; Oxaprotiline Hydrochloride; Oxypertine;  
Paroxetine; Phenelzine Sulfate; Pirandamine Hydrochloride; Pizotiline; Pridefine  
Hydrochloride; Prolintane Hydrochloride; Protriptyline Hydrochloride; Quipazine  
Maleate; Rolicyprine; Seproxetine Hydrochloride; Sertraline Hydrochloride;  
Sibutramine Hydrochloride; Sulpiride; Suritazole; Tametraline Hydrochloride;  
20 Tampramine Fumarate; Tandamine Hydrochloride; Thiazesim Hydrochloride;  
Thozalinone; Tomoxetine Hydrochloride; Trazodone Hydrochloride;  
Trebenzomine Hydrochloride; Trimipramine; Trimipramine Maleate; Venlafaxine

Hydrochloride; Viloxazine Hydrochloride; Zimeldine Hydrochloride;  
Zometapine.

Antihypertensive: Alfuzosin Hydrochloride; Alipamide; Althiazide;  
Amiquinsin Hydrochloride; Amlodipine Besylate; Amlodipine Maleate; Anaritide  
5 Acetate; Atiprosin Maleate; Belfosdil; Bemitradine; Bendacalol Mesylate;  
Bendroflumethiazide; Benzthiazide; Betaxolol Hydrochloride; Bethanidine  
Sulfate; Bevantolol Hydrochloride; Biclodil Hydrochloride; Bisoprolol;  
Bisoprolol Fumarate; Bucindolol Hydrochloride; Bupicomide; Buthiazide;  
Candoxatril; Candoxatrilat; Captopril; Carvedilol; Ceronapril; Chlorothiazide  
10 Sodium; Cicletanine; Cilazapril; Clonidine; Clonidine Hydrochloride; Clopamide;  
Cyclopenthiazide; Cyclothiazide; Darodipine; Debrisoquin Sulfate; Delapril  
Hydrochloride; Diapamide; Diazoxide; Dilevalol Hydrochloride; Diltiazem  
Malate; Ditekiren; Doxazosin Mesylate; Ecadotril; Enalapril Maleate; Enalaprilat;  
Enalkiren; Endralazine Mesylate; Epithiazide; Eprosartan; Eprosartan Mesylate;  
15 Fenoldopam Mesylate; Flavodilol Maleate; Flordipine; Flosequinan; Fosinopril  
Sodium; Fosinoprilat; Guanabenz; Guanabenz Acetate; Guanacline Sulfate;  
Guanadrel Sulfate; Guancydine; Guanethidine Monosulfate; Guanethidine  
Sulfate; Guanfacine Hydrochloride; Guanisoquin Sulfate; Guanoclor Sulfate;  
Guanoctine Hydrochloride; Guanoxabenz; Guanoxan Sulfate; Guanoxyfen  
20 Sulfate; Hydralazine Hydrochloride; Hydralazine Polistirex; Hydroflumethiazide;  
Indacrinone; Indapamide; Indolapril Hydrochloride; Indoramin; Indoramin  
Hydrochloride; Indorenate Hydrochloride; Lacidipine; Leniquinsin;  
Levcromakalim; Lisinopril; Lofexidine Hydrochloride; Losartan Potassium;



Losulazine Hydrochloride; Mebutamate; Mecamylamine Hydrochloride;  
 Medroxalol; Medroxalol Hydrochloride; Methalthiazide; Methyclothiazide;  
 Methyldopa; Methyldopate Hydrochloride; Metipranolol; Metolazone;  
 Metoprolol Fumarate; Metoprolol Succinate; Metyrosine; Minoxidil ; Monatepil  
 5 Maleate; Muzolimine; Nebivolol; Nitrendipine; Ofornine; Pargyline  
 Hydrochloride; Pazoxide; Pelanserin Hydrochloride; Perindopril Erbumine;  
 Phenoxybenzamine Hydrochloride; Pinacidil; Pivopril; Polythiazide; Prazosin  
 Hydrochloride; Primidolol; Prizidilol Hydrochloride; Quinapril Hydrochloride;  
 Quinaprilat; Quinazosin Hydrochloride; Quinelorane Hydrochloride; Quinpirole  
 10 Hydrochloride; Quinuclium Bromide; Ramipril; Rauwolfia Serpentina; Reserpine;  
 Sapisartan Potassium; Saralasin Acetate; Sodium Nitroprusside; Sulfinalol  
 Hydrochloride; Tasosartan; Teludipine Hydrochloride; Temocapril  
 Hydrochloride; Terazosin Hydrochloride; Terlakiren; Tiamenidine; Tiamenidine  
 Hydrochloride; Ticrynafen; Tinabinal; Tiodazosin; Tipentosin Hydrochloride;  
 15 Trichlormethiazide; Trimazosin Hydrochloride; Trimethaphan Camsylate;  
 Trimoxamine Hydrochloride; Tripamide; Xipamide; Zankiren Hydrochloride;  
 Zofenoprilat Arginine.

Anti-inflammatory: Alclofenac; Alclometasone Dipropionate; Algestone  
 Acetonide; Alpha Amylase; Amcinafal; Amcinafide; Amfenac Sodium;  
 20 Amiprilose Hydrochloride; Anakinra; Aniolac; Anitrazafen; Apazone;  
 Balsalazide Disodium; Bendazac; Benoxaprofen; Benzydamine Hydrochloride;  
 Bromelains; Broperamole; Budesonide; Carprofen; Cicloprofen; Cintazone;  
 Cliprofen; Clobetasol Propionate; Clobetasone Butyrate; Clopirac; Cloticasone

Propionate; Cormethasone Acetate; Cortodoxone; Deflazacort; Desonide;  
Desoximetasone; Dexamethasone Dipropionate; Diclofenac Potassium;  
Diclofenac Sodium; Diflorasone Diacetate; Diflumidone Sodium; Diflunisal;  
Difluprednate; Diftalone; Dimethyl Sulfoxide; Drocinnonide; Endrysone;  
5 Enlimomab; Enolicam Sodium; Epirizole; Etodolac; Etofenamate; Felbinac;  
Fenamole; Fenbufen; Fenclofenac; Fenclorac; Fendosal; Fenpipalone; Fentiazac;  
Flazalone; Fluazacort; Flufenamic Acid; Flumizole; Flunisolide Acetate;  
Flunixin; Flunixin Meglumine; Fluocortin Butyl; Fluorometholone Acetate;  
Fluquazone; Flurbiprofen; Fluretofen; Fluticasone Propionate; Furaprofen;  
10 Furobufen; Halcinonide; Halobetasol Propionate; Halopredone Acetate; Ibufenac;  
Ibuprofen; Ibuprofen Aluminum; Ibuprofen Piconol; Ilonidap; Indomethacin;  
Indomethacin Sodium; Indoprofen; Indoxole; Intrazole; Isoflupredone Acetate;  
Isoxepac; Isoxicam; Ketoprofen; Lofemizole Hydrochloride; Lornoxicam;  
Loteprednol Etabonate; Meclofenamate Sodium; Meclofenamic Acid;  
15 Meclorisone Dibutyrate; Mefenamic Acid; Mesalamine; Meseclazone;  
Methylprednisolone Suleptanate; Momiflumate; Nabumetone; Naproxen;  
Naproxen Sodium; Naproxol; Nimazone; Olsalazine Sodium; Orgotein;  
Orpanoxin; Oxaprozin; Oxyphenbutazone; Paranyline Hydrochloride; Pentosan  
Polysulfate Sodium; Phenbutazone Sodium Glycerate; Pirfenidone; Piroxicam;  
20 Piroxicam Cinnamate; Piroxicam Olamine; Pirprofen; Prednazate; Prifelone;  
Prodolic Acid; Proquazone; Proxazole; Proxazole Citrate; Rimexolone;  
Romazarit; Salcolex; Salnacedin; Salsalate; Sanguinarium Chloride; Seclazone;  
Sermetacin; Sudoxicam; Sulindac; Suprofen; Talmetacin; Talniflumate;

Talosalate; Tebufelone; Tenidap; Tenidap Sodium; Tenoxicam; Tesicam;  
Tesimide; Tetrydamine; Tiopinac; Tixocortol Pivalate; Tolmetin; Tolmetin  
Sodium; Triclonide; Triflumidate; Zidometacin; Zomepirac Sodium.

Antinauseant: Buclizine Hydrochloride; Cyclizine Lactate; Naboctate  
5 Hydrochloride .

Antineutropenic: Filgrastim; Lenograstim; Molgramostim; Regramostim;  
Sargramostim.

Antiobsessional agent: Fluvoxamine Maleate.

Antiparkinsonian: Benztropine Mesylate; Biperiden; Biperiden  
10 Hydrochloride; Biperiden Lactate; Carmantadine; Ciladopa Hydrochloride;  
Dopamantine; Ethopropazine Hydrochloride; Lazabemide; Levodopa;  
Lometraline Hydrochloride; Mofegiline Hydrochloride; Naxagolide  
Hydrochloride; Pareptide Sulfate; Procyclidine Hydrochloride; Quinetorane  
Hydrochloride; Ropinirole Hydrochloride; Selegiline Hydrochloride; Tolcapone;  
15 Trihexyphenidyl Hydrochloride. Antiperistaltic: Difenoximide Hydrochloride;  
Difenoxin; Diphenoxylate Hydrochloride; Fluperamide; Lidamidine  
Hydrochloride; Loperamide Hydrochloride; Malethamer; Nufenoxole; Paregoric.

Antipsychotic: Acetophenazine Maleate; Alentemol Hydrobromide;  
Alpertine; Azaperone; Batelapine Maleate; Benperidol; Benzindopyrine  
20 Hydrochloride; Brofbxine; Bromperidol; Bromperidol Decanoate; Butaclamol  
Hydrochloride; Butaperazine; Butaperazine Maleate; Carphenazine Maleate;  
Carvotroline Hydrochloride; Chlorpromazine; Chlorpromazine Hydrochloride;  
Chlorprothixene; Cinperene; Cintriamide; Clomacran Phosphate; Clopenthixol;

- Clopipezide; Clopipezan Mesylate; Cloroperone Hydrochloride; Clothiapine;  
Clothixamide Maleate; Clozapine; Cyclophenazine Hydrochloride; Droperidol;  
Etazolate Hydrochloride; Fenimide; Flucindole; Flumezapine; Fluphenazine  
Decanoate; Fluphenazine Enanthate; Fluphenazine Hydrochloride; Fluspiperone;  
5 Fluspirilene; Flutroline; Gevotroline Hydrochloride; Halopemide; Haloperidol;  
Haloperidol Decanoate; Iloperidone; Imidoline Hydrochloride; Lenperone;  
Mazapertine Succinate; Mesoridazine; Mesoridazine Besylate; Metiapine;  
Milenperone; Milipertine; Molindone Hydrochloride; Naranol Hydrochloride;  
Neflumozide Hydrochloride; Ocaperidone; Olanzapine; Oxiperomide;  
10 Penfluridol; Pentiapine Maleate; Perphenazine; Pimozide; Pinoxepin  
Hydrochloride; Pipamperone; Piperacetazine; Pipotiazine Palmitate; Piquindone  
Hydrochloride; Prochlorperazine Edisylate; Prochlorperazine Maleate; Promazine  
Hydrochloride; Remoxipride; Remoxipride Hydrochloride; Rimcazone  
Hydrochloride; Seperidol Hydrochloride; Sertindole; Setoperone; Spiperone;  
15 Thioridazine; Thioridazine Hydrochloride; Thiothixene; Thiothixene  
Hydrochloride; Tioperidone Hydrochloride; Tiospirone Hydrochloride;  
Trifluoperazine Hydrochloride; Trifluoperidol; Trifluopromazine; Trifluopromazine  
Hydrochloride; Ziprasidone Hydrochloride.
- Appetite suppressant: Dexfenfluramine Hydrochloride; Phendimetrazine  
20 Tartrate; Phentermine Hydrochloride.
- Blood glucose regulators: Human insulin; Glucagon; Tolazamide;  
Tolbutamide; Chlorpropamide; Acetohexamide and Glipizide.

Carbonic anhydrase inhibitor: Acetazolamide; Acetazolamide Sodium, Dichlorphenamide; Dorzolamide Hydrochloride; Methazolamide; Sezolamide Hydrochloride.

Cardiac depressant: Acecainide Hydrochloride; Acetylcholine Chloride;  
5 Actisomide; Adenosine; Amiodarone; Aprindine; Aprindine Hydrochloride;  
Artilide Fumarate; Azimilide Dihydrochloride; Bidisomide; Bucainide Maleate;  
Bucromarone; Butopropine Hydrochloride; Capobenat Sodium; Capobenat Acid;  
Cifenline; Cifenline Succinate; Clofilium Phosphate; Disobutamide;  
Disopyramide; Disopyramide Phosphate; Dofetilide; Drobuline; Edifolone  
10 Acetate; Emilium Tosylate; Encainide Hydrochloride; Flecainide Acetate;  
Ibutilide Fumarate; Indecainide Hydrochloride; Ipazilide Fumarate; Lorajmine  
Hydrochloride; Lorcanide Hydrochloride; Meobentine Sulfate; Mexiletine  
Hydrochloride; Modecainide; Moricizine; Oxiramide; Pirmenol Hydrochloride;  
Pirolazamide; Pranolium Chloride; Procainamide Hydrochloride; Propafenone  
15 Hydrochloride; Pyrinoline; Quindonium Bromide; Quinidine Gluconate;  
Quinidine Sulfate; Recainam Hydrochloride; Recainam Tosylate; Risotilide  
Hydrochloride; Ropitoin Hydrochloride; Sematilide Hydrochloride; Suricainide  
Maleate; Tocainide; Tocainide Hydrochloride; Transcainide.

Cardiotonic: Actodigin; Amrinone; Bemoradan; Butopamine; Carbazeran;  
20 Carsatrin Succinate; Deslanoside; Digitalis; Digitoxin; Digoxin; Dobutamine;  
Dobutamine Hydrochloride; Dobutamine Lactobionate; Dobutamine Tartrate;  
Enoximone; Imazodan Hydrochloride; Indolidan; Isomazole Hydrochloride;  
Levdo butamine Lactobionate; Lixazinone Sulfate; Medorinone; Milrinone;

Pelrinone Hydrochloride; Pimobendan; Piroximone; Prinoxodan; Proscillaridin;  
Quazinone; Tazolol Hydrochloride; Vesnarinone.

Cardiovascular agent: Dopexamine; Dopexamine Hydrochloride.

Choleretic: Dehydrocholic Acid; Fencibutirol; Hymecromone; Piprozolin;

5 Sincalide; Tocamphyl.

Cholinergic: Aceclidine; Bethanechol Chloride; Carbachol; Demecarium  
Bromide; Dexpanthenol; Echothiophate Iodide; Isoflurophate; Methacholine  
Chloride; Neostigmine Bromide; Neostigmine Methylsulfate; Physostigmine;  
Physostigmine Salicylate; Physostigmine Sulfate; Pilocarpine; Pilocarpine  
10 Hydrochloride; Pilocarpine Nitrate; Pyridostigmine Bromide.

Cholinergic agonist: Xanomeline; Xanomeline Tartrate.

Cholinesterase Deactivator: Obidoxime Chloride; Pralidoxime Chloride;  
Pralidoxime Iodide; Pralidoxime Mesylate.

Coccidiostat: Arprinocid; Narasin; Semduramicin; Semduramicin Sodium.

15 Cognition adjuvant: Ergoloid Mesylates; Piracetam; Pramiracetam  
Hydrochloride; Pramiracetam Sulfate; Tacrine Hydrochloride.

Cognition enhancer: Besipirdine Hydrochloride; Linopirdine; Sibopirdine.

Hormone: Diethylstilbestrol; Progesterone; 17 hydroxy progesterone;  
Medroxyprogesterone; Norgestrel; Norethynodrel; Estradiol; Megestrol  
20 (Megace); Norethindrone; Levonorgestrel; Ethyndiol; Ethinyl estradiol;  
Mestranol; Estrone; Equilin; 17 alpha dihydroequilin; equilenin; 17 alpha  
dihydroequilenin; 17 alpha estradiol; 17 beta estradiol; Leuprolide (lupron);  
Glucagon; Testolactone; Clomiphene; Human menopausal gonadotropins; Human

- chorionic gonadotropin; Urofollitropin; Bromocriptine; Gonadorelin; Luteinizing hormone releasing hormone and analogs; Gonadotropins; Danazol; Testosterone; Dehydroepiandrosterone; Androstenedione; Dihydroestosterone; Relaxin; Oxytocin; Vasopressin; Folliculostatin; Follicle regulatory protein;
- 5 Gonadotrinins; Oocyte maturation inhibitor; Insulin growth factor; Follicle Stimulating Hormone; Luteinizing hormone; Tamoxifen.; Corticorelin Ovine Triflutate; Cosyntropin; Metogest; Pituitary, Posterior; Seractide Acetate; Somalapor; Somatrem; Somatropin; Somenopor; Somidobove.
- Memory adjuvant: Dimoxamine Hydrochloride; Ribaminol.
- 10 Mental performance enhancer: Aniracetam.
- Mood regulator: Fengabine.
- Neuroleptic: Duoperone Fumarate; Risperidone.
- Neuroprotective: Dizocilpine Maleate.
- Psychotropic: Minaprine.
- 15 Relaxant: Adiphenine Hydrochloride; Alcuronium Chloride; Aminophylline; Azumolene Sodium; Baclofen; Benzoctamine Hydrochloride; Carisoprodol; Chlorphenesin Carbamate; Chlorzoxazone; Cinflumide; Cinnamedrine; Clodanolene; Cyclobenzaprine Hydrochloride; Dantrolene; Dantrolene Sodium; Fenalamide; Fenyripol Hydrochloride; Fetoxylyate
- 20 Hydrochloride; Flavoxate Hydrochloride; Fletazepam; Flumetramide;-Flurazepam Hydrochloride; Hexafluorenium Bromide; Isomylamine Hydrochloride; Lorbamate; Mebeverine Hydrochloride; Mesuprine Hydrochloride; Metaxalone; Methocarbamol; Methixene Hydrochloride; Nafomine Malate; Nelezaprine

Maleate; Papaverine Hydrochloride; Pipoxolan Hydrochloride; Quinctolate;  
Ritodrine; Ritodrine Hydrochloride; Rolodine; Theophylline Sodium Glycinate;  
Thiphenamil Hydrochloride; Xilobam.

Sedative-hypnotic: Allobarbital; Alonimid; Alprazolam; Amobarbital  
5 Sodium; Bentazepam; Brotizolam; Butabarbital; Butabarbital Sodium; Butalbital;  
Capuride; Carbocloral; Chloral Betaine; Chloral Hydrate; Chlordiazepoxide  
Hydrochloride; Cloperidone Hydrochloride; Clorethate; Cyprazepam; Dexclamol  
Hydrochloride; Diazepam; Dichloralphenazone; Estazolam; Ethchlorvynol;  
Etomidate; Fenobam; Flunitrazepam; Fosazepam; Glutethimide; Halazepam;  
10 Lormetazepam; Mecloqualone; Meprobamate; Methaqualone; Midaflur;  
Paraldehyde; Pentobarbital; Pentobarbital Sodium; Perlapine; Prazepam;  
Quazepam; Reclazepam; Roletamide; Secobarbital; Secobarbital Sodium;  
Suproclonine; Thalidomide; Tracazolate; Trepipam Maleate; Triazolam;  
Tricetamide; Triclofos Sodium; Trimetozine; Uldazepam; Zaleplon; Zolazepam  
15 Hydrochloride; Zolpidem Tartrate.

Serotonin antagonist: Altanserin Tartrate; Amesergide; Ketanserin;  
Ritanserin.

Serotonin inhibitor: Cinanserin Hydrochloride; Fenclonine; Fonazine  
Mesylate; Xylamidine Tosylate.

20 Serotonin receptor antagonist: Tropanserin Hydrochloride.

Stimulant: Amfonelic Acid; Amphetamine Sulfate; Ampyzine Sulfate;  
Arbutamine Hydrochloride; Azabon; Caffeine; Ceruletide; Ceruletide  
Diethylamine; Cisapride; Dazopride Fumarate; Dextroamphetamine;



Dextroamphetamine Sulfate; Difluanine Hydrochloride; Dimeflin  
Hydrochloride; Doxapram Hydrochloride; Etryptamine Acetate; Ethamivan;  
Fenethylamine Hydrochloride; Flubanilate Hydrochloride; Flurothyl; Histamine  
Phosphate; Indriline Hydrochloride; Mefexamide; Methamphetamine Hydrochloride;  
5 ride; Methylphenidate Hydrochloride; Pemoline; Pyrovalerone Hydrochloride;  
Xamoterol; Xamoterol Fumarate.

Synergist: Proadifen Hydrochloride.

Thyroid hormone: Levothyroxine Sodium; Liothyronine Sodium; Liotrix.

Thyroid inhibitor: Methimazole; Propylthiouracil.

10 Thyromimetic: Thyromedan Hydrochloride.

Cerebral ischemia agents: Dextrophan Hydrochloride.

Vasoconstrictor: Angiotensin Amide; Felypressin; Methysergide;  
Methysergide Maleate.

Vasodilator: Alprostadil; Azaclozine Hydrochloride; Bamethan Sulfate;  
15 Bepridil Hydrochloride; Buterizine; Cetiedil Citrate; Chromonar Hydrochloride;  
Clonitrate; Diltiazem Hydrochloride; Dipyridamole; Droprenilamine; Erythrityl  
Tetranitrate; Felodipine; Flunarizine Hydrochloride; Fostedil; Hexobendine;  
Inositol Niacinate; Iproxamine Hydrochloride; Isosorbide Dinitrate; Isosorbide  
Mononitrate; Isoxsuprine Hydrochloride; Lidoflazine; Mefenidil; Mefenidil  
20 Fumarate; Mibefradil Dihydrochloride; Mioflazine Hydrochloride; Mixidine;  
Nafronyl Oxalate; Nicardipine Hydrochloride; Nicergoline; Nicorandil; Nicotinyl  
Alcohol; Nifedipine; Nimodipine; Nisoldipine; Oxfenicine; Oxprenolol  
Hydrochloride; Pentaerythritol Tetranitrate; Pentoxifylline; Pentritinol;

Perhexiline Maleate; Pindolol; Pirsidomine; Prenylamine; Propatyl Nitrate; Suloctidil; Terodiline Hydrochloride; Tipropidil Hydrochloride; Tolazoline Hydrochloride; Xanthinol Niacinate.

5           Specifically, topiramate may be administered in combination with other medications to treat certain symptoms and disorders including:

- I. Treatment of Binge Eating (Binge Eating Disorder, Bulimia Nervosa, Anorexia Nervosa with Binge eating) with serotonin re-uptake inhibitors (*e.g.*, citalopram (CELEXA), clomipramine (ANAFRANIL)), fluoxetine (PROZAC),  
10       fluvoxamine (LUVOX), venlafaxine (EFFEXOR), other antidepressants (*e.g.*, bupropion (WELLBUTRIN) nefazodone (SERZONE), tricyclics (*e.g.*, NORPRAMIN and PAMELOR), trazodone (DESYREL), Substance P antagonists), psychostimulants, (*e.g.*, d-amphetamine, phentermine; and sibutramine (MERIDIA)) and orlistat.
- 15       II. Treatment of overweight/obesity condition with sibutramine (MERIDIA); psychostimulants, (*e.g.*, d-amphetamine, phentermine) and orlistat.
- III. Treatment of nicotine addiction/smoking cessation with bupropion (ZYBAN), serotonin reuptake inhibitors, nicotine patches and gum, and other antidepressants.
- 20       IV. Treatment of alcohol abuse/dependence (alcoholism) with naltrexone (REVIA), serotonin reuptake inhibitors, and other antidepressants.
- V. Treatment of other impulse control disorders (behavioral addictions) with serotonin reuptake inhibitors, lithium, valproic acid or divalproex sodium (*e.g.*,

DEPAKENE or DEPAKOTE), other antidepressants, naltrexone, atypical antipsychotics, (*e.g.*, olanzapine (ZYPREXA), quetiapine (SEROQUEL), risperidone (RISPERDAL), ziprasidone) and other mood stabilizers (*e.g.*, carbamazepine)

- 5 VI. Treatment of paraphilias/sexual addictions with serotonin reuptake inhibitors, lithium, divalproex sodium/valproic acid, antiandrogen medications (*e.g.*, medroxyprogesterone, gonadotropin-releasing hormone (GnRH) agonists), other antidepressants, and other mood stabilizers (*e.g.*, carbamazepine).

- 10 When administered, the formulations of the invention are applied in pharmaceutically acceptable amounts and in pharmaceutically acceptable compositions. Such preparations may routinely contain salts, buffering agents, preservatives, compatible carriers, and optionally other therapeutic ingredients. When used in medicine the salts should be pharmaceutically acceptable, but non-pharmaceutically acceptable salts may
- 15 conveniently be used to prepare pharmaceutically acceptable salts thereof and are not excluded from the scope of the invention. Such pharmacologically and pharmaceutically acceptable salts include, but are not limited to, those prepared from the following acids: hydrochloric, hydrobromic, sulfuric, nitric, phosphoric, maleic, acetic, salicylic, p-toluene sulfonic, tartaric, citric, methane sulfonic, formic, malonic, succinic, naphthalene-2-
- 20 sulfonic, and benzene sulfonic. Also, pharmaceutically acceptable salts can be prepared as alkaline metal or alkaline earth salts, such as sodium, potassium or calcium salts.

Suitable buffering agents include: acetic acid and a salt (1-2% W/V); citric acid and a salt (1-3% W/V); boric acid and a salt (0.5-2.5% W/V); and phosphoric acid and a

salt (0.8-2% W/V). Suitable preservatives include benzalkonium chloride (0.003-0.03% W/V); chlorobutanol (0.3-0.9% W/V); parabens (0.01-0.25% W/V) and thimerosal (0.004-0.02% W/V).

In the present invention, the sulfamide derivatives are administered in safe and effective amounts. An effective amount means that amount necessary to delay the onset of, inhibit the progression of, halt altogether the onset or progression of or diagnose the particular condition being treated. In general, an effective amount for treating an ICD will be that amount necessary to inhibit mammalian symptoms of the particular ICD in-situ. When administered to a subject, effective amounts will depend, of course, on the particular condition being treated; the severity of the condition; individual patient parameters including age, physical condition, size and weight; concurrent treatment; frequency of treatment; and the mode of administration. These factors are well known to those of ordinary skill in the art and can be addressed with no more than routine experimentation. It is preferred generally that a minimum dose be used, that is, the lowest safe dosage that provides appropriate relief of symptoms.

Dosage may be adjusted appropriately to achieve desired drug levels, locally or systemically. Generally, daily oral doses of active compounds will be from about 0.01 mg/kg per day to 2000 mg/kg per day. It is expected that IV doses in the range of about 1 to 1000 mg/cm<sup>3</sup> per day will be effective. In the event that the response in a subject is insufficient at such doses, even higher doses (or effective higher doses by a different, more localized delivery route) may be employed to the extent that patient tolerance permits. Continuous IV dosing over, for example 24 hours or multiple doses per day is contemplated to achieve appropriate systemic levels of compounds.

A variety of administration routes are available. The particular mode selected will depend of course, upon the particular drug selected, the severity of the disease state(s) being treated and the dosage required for therapeutic efficacy. The methods of this invention, generally speaking, may be practiced using any mode of administration that is medically acceptable, meaning any mode that produces effective levels of the active compounds without causing clinically unacceptable adverse effects. Such modes of administration include oral, rectal, sublingual, topical, nasal, transdermal or parenteral routes. The term "parenteral" includes subcutaneous, intravenous, intramuscular, or infusion. Intravenous routes are preferred.

10       The compositions may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. In general, the compositions are prepared by uniformly and intimately bringing the compounds into association with a liquid carrier, a finely divided solid carrier, or both, and then, if necessary, shaping the product.

15       Compositions suitable for oral administration may be presented as discrete units such as capsules, cachets, tablets, or lozenges, each containing a predetermined amount of the active compound. Other compositions include suspensions in aqueous liquors or non-aqueous liquids such as a syrup, an elixir, or an emulsion.

Other delivery systems can include time-release, delayed release or sustained release delivery systems. Such systems can avoid repeated administrations of the active compounds of the invention, increasing convenience to the subject and the physician. Many types of release delivery systems are available and known to those of ordinary skill in the art. They include polymer based systems such as polylactic and polyglycolic acid,

polyanhydrides and polycaprolactone; nonpolymer systems that are lipids including sterols such as cholesterol, cholesterol esters and fatty acids or neutral fats such as mono-, di and triglycerides; hydrogel release systems; silastic systems; peptide based systems; wax coatings, compressed tablets using conventional binders and excipients, partially  
5 fused implants and the like. In addition, a pump-based hardware delivery system can be used, some of which are adapted for implantation.

A long-term sustained release implant also may be used. "Long-term" release, as used herein, means that the implant is constructed and arranged to deliver therapeutic levels of the active ingredient for at least 30 days, and preferably 60 days. Long-term  
10 sustained release implants are well known to those of ordinary skill in the art and include some of the release systems described above.

#### **Examples.**

In the examples, patients were treated with open-label topiramate starting at 25 mg/qHS  
15 and the dosage increased by the patient in 25 mg increments as tolerated by the subjects until a response is seen up to a maximum of 1200 mg.

Table 1. Patients with Binge Eating Disorder (BED) Treated Clinically with Open-Label Topiramate (as of 12/16/98)

Pt #	Pt ID	Reasons Topiramate Begun	Max Dose (mg/day)	Response
1	LJM	BD (BED) (Obesity)	1200	Remission of BD Remission of BED Loss of 107 lbs.
2	CEJ	BD BED Overweight	150	Mild improvement of BD, Moderate decrease of BED, Loss of 5 lbs. Discontinued due to sedation, cognitive dulling
3	JAC	BD BED Obesity	1200	Moderate improvement of BD. Remission of BED Loss of 50.5 lbs
4	JEB	BD BED Overweight	900	Moderate improvement of BD, Marked improvement of BED, Loss of 30.5 lbs.
5	KCW	BED Obesity Compulsive Buying (BD, in remission)	100  100	First trial: No response of BED, discontinued due to GI distress  Second Trial: Worsening of BD Remission of BED Loss of 11 lbs. Remission of Compulsive Buying
6	JB	BED Overweight	100	No response of BED No weight change

- 5 Key: BD=Bipolar Disorder; BED=Binge Eating Disorder; Pt.=patient; D/C= topiramate treatment discontinued; Cont.= topiramate treatment continued; GI=gastrointestinal

Table 2. Patients with Overeating, Overweight, and Obesity Treated Clinically with Topiramate (as of 12/16/98)

Pt. #	Pt. ID	Reasons Topiramate Begun	Max dose (mg/day)	Response
1	BAA	BD Obesity (BED, in remission)	700	No response of BD Loss of 9lbs. (293-284)
2	HTB	BD Overeating Overweight	300	Remission of BD Remission of overeating Loss of 16 lbs. (248-232) Discontinued 2° illness in remission
3	ADJ	BD Overeating, Overweight	400	Moderate improvement of BD Moderate improvement of overeating Loss of 7 lbs. (239-232)
4	TK	BD Overeating Overweight	350	No response of BD Mild improvement in overeating Loss of 5 lbs. (238-233)
5	JCJ	BD Overeating Overweight	1. 250 2. 200	First Trial: Worsening of BD Mild decrease of overeating Loss of 7 lbs. (152-145) Second Trial Mild improvement of BD Mild decrease of overeating Gain of 1lb. (154-155 lbs.)
6	KDC	Overeating Overweight (BD, in remission)	800	Worsening of BD Loss of 9 lbs. (208-199lbs)
7	NLR	BD Overweight	300	Worsening of BD Loss of 46 lbs. (196-150 lbs.) Discontinued due to anorexia
8	PR	BD Overeating Overweight	700	Moderate improvement of BD Marked improvement in overeating Loss of 19 lbs. (185-166) Discontinued due to G.I. Distress


Key: Pt.=patient; BD=Bipolar Disorder; BED=Binge Eating Disorder; D/C=topiramate treatment discontinued; cont=topiramate treatment continued; GI=gastrointestinal



**CLAIMS**


What is claimed is:

- 1) A method for treating an Impulse Control Disorder comprising administering to a mammal afflicted with such condition a therapeutically effective amount for treating such condition of a compound of the formula I:



wherein

- X is CH<sub>2</sub> or oxygen;
- R<sub>1</sub> is hydrogen or alkyl; and
- R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub> and R<sub>5</sub> are independently hydrogen or lower alkyl and, when X is CH<sub>2</sub>, R<sub>4</sub> and R<sub>5</sub> may be alkene groups joined to form a benzene ring and, when X is oxygen, R<sub>2</sub> and R<sub>3</sub> and/or R<sub>4</sub> and R<sub>5</sub> together may be a methylenedioxy group of the following formula (II):



wherein

R<sub>6</sub> and R<sub>7</sub> are the same or different and are hydrogen, lower alkyl or are alkyl and are joined to form a cyclopentyl or cyclohexyl ring.

- 2) The method of claim 1 wherein the compound is selected from the group consisting of 2,3:4,5-bis-O-(1-methylethylidene)- beta -D-fructopyranose sulfamate, 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose sulfamate, 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -L-fructopyranose sulfamate, 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose methylsulfamate, 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose butylsulfamate, 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose ethylsulfamate, 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose octylsulfamate, 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose 2-propenylsulfamate, 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose phenylmethylsulfamate, 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose cyclopropylsulfamate, 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose cyclobutylsulfamate, 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose (2,2,2-trifluoroethyl)sulfamate, 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose dimethylsulfamate, 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose diethylsulfamate, 2,3-O-(1-methylethylidene)-4,5-O-sulfonyl- beta -D-fructopyranose azidosulfate, (S)-2,3-O-(1-methylethylidene)-4,5-O-sulfinyl- beta -D-fructopyranose sulfamate, (R)-2,3-O-(1-methylethylidene)-4,5-O-sulfinyl- beta -D-fructopyranose sulfamate, 2,3-O-(1-ethylpropylidene)-4,5-O-sulfonyl- beta -D-fructopyranose sulfamate, 2,3-O-(1-methylethylidene)-4,5-O-[N-(4-methylbenzenesulfonyl)imidodisulfinyl]- beta -D-fructopyranose sulfamate, 2,3-O-(1-methylethylidene)-4,5-O-[N-(4-

methylbenzenesulfonyl)imidodisulfonyl]- beta -D-fructopyranose sulfamate, 2,3-O-(cyclohexylidene)-4,5-O-sulfonyl- beta -D-fructopyranose sulfamate, (S)-4,5-O-[N-(1,1-dimethylethoxycarbonyl)imidodisulfonyl]-2,3-O-(1-methylethylidene)- beta -D-fructopyranose sulfamate, and the pharmaceutically acceptable salts thereof.

- 3) The method of claim 1 wherein the compound is selected from the group consisting of (1R,2R,3S,4S)-(1,2:3,4-di-O-methylethylidenecyclohexan-1,2,3,4-tetraol-4-yl)methyl sulfamate and (1R,2S,3S,4S)-(3,4-O-methylethylidene-1,2-O-sulfinylcyclohexan-1,2,3,4-tetraol-4-yl)methyl sulfamate.
- 4) The method of claim 1 wherein the compound of formula I is topiramate.
- 5) The method of claim 1, wherein the therapeutically effective amount is of from about 15 to about 2000 mg per day.
- 6) The method of claim 1, wherein the therapeutically effective amount is of from about 25 to about 750 mg per day.

- 7) The method of claim 1 wherein the Impulse Control Disorder is selected from the group consisting of intermittent explosive disorder (IED), kleptomania, pathological gambling, pyromania, trichotillomania, compulsive buying or shopping, repetitive self-mutilation, nonparaphilic sexual addictions, severe nail biting, compulsive skin picking, personality disorders with impulsive features, attention deficit/hyperactivity disorder, eating disorders characterized by binge eating, and substance use disorders.
- 8) The method of claim 1 wherein the compound is used in conjunction with one or more other drug compounds selected from the group consisting of adrenergics, adrenocortical steroids, adrenocortical suppressants, aldosterone antagonists, amino acids, analeptics, analgesics, anorectic compounds, anorexics, anti-anxiety agents, antidepressants, antihypertensives, anti-inflammatorys, antinauseants, antineutropenics, antiobsessional agents, antiparkinsonians, antipsychotics, appetite suppressants, blood glucose regulators, carbonic anhydrase inhibitors, cardiotonics, cardiovascular agents, choleretics, cholinergics, cholinergic agonists, cholinesterase deactivators, cognition adjuvants, cognition enhancers, hormones, memory adjuvants, mental performance enhancers, mood regulators, neuroleptics, neuroprotectives, psychotropics, relaxants, sedative-hypnotics, serotonin antagonists, serotonin inhibitors, serotonin receptor antagonists, stimulants, thyroid hormones, thyroid inhibitors, thyromimetics, cerebral ischemia agents, vasoconstrictors, and vasodilators.

- 9) The method of claim 1 wherein the Impulse Control Disorder is an eating disorder and the compound is used in conjunction with one or more other drug compounds selected from the group consisting of serotonin re-uptake inhibitors, antidepressants, psychostimulants, and orlistat.
- 10) The method of claim 1 wherein the Impulse Control Disorder is an overweight/obesity condition and the compound is used in conjunction with one or more other drug compounds selected from the group consisting of sibutramine, psychostimulants, and orlistat.
- 11) The method of claim 1 wherein the Impulse Control Disorder is a nicotine addiction condition and the compound is used in conjunction with one or more other drug compounds selected from the group consisting of bupropion, serotonin reuptake inhibitors, nicotine, and antidepressants.
- 12) The method of claim 1 wherein the Impulse Control Disorder is an alcohol abuse/dependence condition and the compound is used in conjunction with one or more other drug compounds selected from the group consisting of naltrexone, serotonin reuptake inhibitors, and other antidepressants.

- 13) The method of claim 1 wherein the Impulse Control Disorder is a behavioral addiction condition and the compound is used in conjunction with one or more other drug compounds selected from the group consisting of serotonin reuptake inhibitors, lithium, valproic acid or divalproex sodium, other antidepressants, 5 naltrexone, atypical antipsychotics, and other mood stabilizers.
- 14) The method of claim 1 wherein the Impulse Control Disorder is a paraphilias/sexual addiction condition and the compound is used in conjunction with one or more other drug compounds selected from the group consisting of serotonin reuptake inhibitors, lithium, divalproex sodium/valproic acid, 5 antiandrogen agents, other antidepressants, and other mood stabilizers.